

# Protection without poison: Why tropical ozone maximizes in the interior of the atmosphere

Aaron Match<sup>1</sup>, Edwin P. Gerber<sup>1</sup>, and Stephan Fueglistaler<sup>2</sup>

<sup>1</sup>Center for Atmosphere Ocean Science, Courant Institute of Mathematical Sciences, New York University, New York, NY, USA

<sup>2</sup>Program in Atmospheric and Oceanic Sciences, and Department of Geosciences, Princeton University, Princeton, NJ, USA


**Correspondence:** Aaron Match (aaron.match@nyu.edu)

**Abstract.** The number density of ozone,  $[O_3]$ , maximizes around 26 km in the tropics, protecting life from harmful ultraviolet light (UV) without poisoning it at the surface. Textbooks explain this interior maximum with two paradigms: (1) the *source-controlled paradigm* explains  $[O_3]$  to maximize where its source maximizes between abundant photons aloft and abundant  $[O_2]$  below, and (2) the *source/sink competition paradigm*, inspired by the Chapman cycle, explains ozone to scale with  $[O_2]$  and the photolytic source/sink ratio. Yet, each paradigm's prediction for the altitude of peak  $[O_3]$  is off by 10 km, reflecting their well-known omission of ozone sinks from catalytic cycles and transport. We present a minimal, steady-state theory for the tropical stratospheric  $[O_3]$  maximum, accurate to within 1 km, formulated in terms of the dominant ozone sinks. These sinks are represented simply by augmenting the Chapman cycle with linear damping of O and  $O_3$ , leading to the Chapman+2 model. The Chapman+2 model correctly simulates peak tropical  $[O_3]$  at 26 km, yet this peak is not explained by either paradigm. Instead, the peak is newly explained by the transition from an O-damped regime aloft to an  $O_3$ -damped regime below. An explicit analytical expression is derived for ozone under gray radiation. This theory accurately predicts an interior maximum of ozone and correctly predicts that an increase in top-of-atmosphere UV leads to a downward shift of peak  $[O_3]$ , due to a downward shift in the regime transition, a result not even qualitatively predicted by the existing paradigms.

## 1 Introduction

Ozone's presence in the stratosphere protects life from harmful UV radiation. It was the absence of this high-energy radiation at the surface that enabled Hartley to deduce the existence of the ozone layer (Hartley, 1881). In addition to protecting life from UV, ozone is also a strong oxidizing agent, making it poisonous to lungs and plant tissues. Thus, by maximizing well above the surface, around 26 km in the tropics, the ozone layer provides protection without poison.

This paper seeks to understand the tropical stratospheric maximum of ozone number density, denoted  $[O_3]$  ( $\text{molec cm}^{-3}$ ). The tropical stratospheric peak in  $[O_3]$  is robust across observational datasets. As an observational benchmark, this paper uses the homogenized satellite dataset SWOOSH (Davis et al., 2016), averaged over the tropics ( $30^\circ\text{S}$ - $30^\circ\text{N}$ ) and from 1984-2023. In SWOOSH, monthly tropical  $[O_3]$  peaks at 26 km, deviating only about 10% of the time up or down from this altitude by at most one vertical level of roughly 1 km. The tropical stratospheric peak in  $[O_3]$  is also robust in state-of-the-art chemistry-climate models, which successfully reproduce this interior maximum (e.g., Keeble et al., 2021). But, since these chemistry-



**Figure 1.** Two textbook paradigms for explaining the interior maximum of ozone. (a) In the source-controlled paradigm, ozone scales with its production rate given by the product of the photolysis rate ( $J_{O_2}$ ), which decreases towards the surface, and the number density of  $[O_2]$ , which increases exponentially towards the surface. Their product maximizes at a sweet spot. (b) In the source/sink competition paradigm, ozone scales as in the Chapman cycle, with dependence on  $[O_2]$  and on the ratio of photolysis rates of  $O_2$  and  $O_3$ . In the source/sink competition paradigm, photolysis of  $O_3$  suppresses  $O_3$ .

25 climate models include a complex representation of the global atmospheric circulation and hundreds of chemical reactions, the reasons for this emergent structure can be hard to discern. Here, we seek to drastically reduce the apparent complexity by distilling the minimal set of physical and chemical processes required to explain this robust feature of observations and models.

Explaining the interior maximum of tropical ozone is an old problem, and the modern theory for the structure of the ozone layer began almost a century ago when Sydney Chapman demonstrated that the ozone layer is formed via ultraviolet (UV) photochemistry (Chapman, 1930). Chapman showed how a motionless atmospheric column illuminated by UV could produce an ozone layer through photochemical cycles of  $O$ ,  $O_2$ , and  $O_3$ . Explaining why ozone has an interior maximum is now a standard part of atmospheric chemistry curricula. We surveyed eight atmospheric radiation and chemistry textbooks that explained why ozone has an interior maximum<sup>1</sup>, and found that, when explaining the structure of the ozone layer, all introduce the Chapman cycle, even as most subsequently note its important omissions of catalytic cycles and transport. What has not  
 30 been previously noted is that, when explaining the interior maximum, textbooks invoke two qualitatively different paradigms, with half of the surveyed textbooks invoking each paradigm.

The first paradigm, is the *source-controlled paradigm*. It asserts that the interior maximum of tropical  $[O_3]$  is dictated by the interior maximum in the ozone production rate,  $J_{O_2}[O_2]$  ( $\text{molec cm}^{-3} \text{ s}^{-1}$ ), where  $J_{O_2}$  is the photolysis rate of  $O_2$  ( $\text{s}^{-1}$ ),

<sup>1</sup>Dobson (1963); Dutsch (1968); Jacob (1999); Liou (2002); McElroy (2002); Kump et al. (2011); Visconti (2016); Seinfeld and Pandis (2016)


and  $[O_2]$  is the number density of  $O_2$  (molec  $cm^{-3}$ ).  $J_{O_2}$  is large aloft but attenuates rapidly towards the surface, whereas  $[O_2]$  increases exponentially towards the surface, so their product maximizes at a “sweet spot” where there are both enough photons and enough  $O_2$  molecules to yield a large  $O_3$  production rate. The sweet spot suggests that the interior maximum of ozone is a fundamental consequence of radiative attenuation through an exponentially-distributed absorber (Jacob, 1999), for which the radiative absorption rate (photons  $cm^{-3} s^{-1}$ ) maximizes where optical depth equals one. A cartoon version of this paradigm is shown in Fig. 1a.

The second paradigm is the *source/sink competition paradigm* (Fig. 1b). The source/sink competition paradigm uses the precise functional form of ozone derived from the Chapman cycle, where the vertical structure of the ozone layer predominantly scales as  $(J_{O_2}/J_{O_3})^{1/2}[O_2]^{3/2}$ . The photolysis rate of  $O_3$ ,  $J_{O_3}$ , enters explicitly into the denominator of this expression because photolysis of ozone contributes to the sink of ozone in the Chapman cycle by liberating atomic oxygen that can then bond with  $O_3$  to destroy it. The source/sink competition paradigm suggests that the interior maximum of tropical  $[O_3]$  arises due to competition between a photolytic source and photolytic sink, playing out within the photochemical context of the Chapman cycle. A cartoon version of the source/sink competition paradigm is shown in Fig. 1b.

Given that these paradigms yield different scaling relationships for the vertical structure of the ozone layer, the question arises: is this a theoretically ambiguous case where different paradigms yield a consistent, accurate prediction? This can be tested by evaluating the paradigmatic scalings, which we calculate based on the photolysis rates that result when incoming UV is attenuated by  $O_2$  and observed  $O_3$ , treating radiative attenuation as described in Section 2. The results are shown in Figure 2. The source-controlled paradigm (blue) predicts  $[O_3]$  to maximize at 38 km, and the source/sink competition paradigm (red) predicts  $[O_3]$  to maximize at 15 km. Thus, these paradigmatic scalings are neither consistent (they disagree with each other by more than three atmospheric scale heights) nor accurate (they each disagree with observations by more than 10 km).

The roots of these inconsistencies and inaccuracies can be hypothesized based on prior knowledge: these paradigms treat ozone sinks inconsistently and inaccurately, and both are known to neglect the dominant ozone sinks. The source-controlled paradigm neglects any structural contribution from the sink, tantamount to assuming a damping-like sink of the form  $\partial[O_3]/\partial t = -\kappa_{O_3}[O_3] + \dots$ , with insignificant vertical structure in  $\kappa_{O_3}$ . The source/sink competition paradigm accounts only for the Chapman cycle sink of ozone from the reaction of O and  $O_3$ , which is known to be minor in observations. Neglected in these paradigms are the dominant observed sinks of ozone: catalytic cycles and transport (e.g., Bates and Nicolet, 1950; Crutzen, 1970; Brasseur and Solomon, 2005).

We seek a minimal, steady-state theory for the tropical stratospheric  $[O_3]$  maximum that invokes realistic sinks from catalytic cycles and transport and yields a prediction for the interior maximum of ozone that is accurate to approximately 1 km. To develop a theory that invokes realistic sinks, we bridge the gap between simple theories and comprehensive simulations by augmenting the Chapman cycle with two linear damping reactions that represent the destruction of either O or  $O_3$  by catalytic cycles and transport. We call our resulting photochemical system the *Chapman+2 model* (sensitivities in which were recently analyzed in Match et al., 2024). The damping coefficients of O and  $O_3$  must be constrained by the known magnitudes of catalytic cycles and transport.



**Figure 2.** The vertical structure of the tropical ozone layer in observations (black) and from the scalings within the two paradigms (blue and red). Observations of  $O_3$  are from homogenized satellite data in the SWOOSH dataset (Davis et al., 2016) averaged from 1984–2023 and over the tropics from  $30^\circ\text{S}$ – $30^\circ\text{N}$ . The photolysis rates  $J_{O_2}$  and  $J_{O_3}$  are calculated based on optical depth-based radiative transfer with overhead sun and absorption by  $O_2$  and  $O_3$  (using the observed profile), otherwise following the methods in Section 2. All profiles are normalized by their maximum value, whose altitude is starred and labeled.

Whether the damping is primarily of  $O$  versus of  $O_3$  turns out to lead to qualitatively different mechanisms for ozone structure, a surprising result given that  $O$  and  $O_3$  are often treated as conceptually fungible within the chemical family of odd oxygen ( $O_x \equiv O + O_3$ ) (Section 2). In the **O-damped regime**, the destruction of ozone is rate-limited by the availability of atomic oxygen, which must be produced through photolysis of ozone. On its own, the O-damped regime produces an interior maximum at the same altitude as predicted by the source/sink competition paradigm. The  **$O_3$ -damped regime** produces an interior maximum at the same altitude as predicted by the source-controlled paradigm.

Today’s tropical stratosphere occupies each regime at different altitudes, with the transition from an O-damped regime aloft to an  $O_3$ -damped regime below at 26 km, co-located with the ozone maximum (Section 3). Although each paradigm is capable of producing an interior maximum of ozone, neither can successfully explain the observed altitude of the tropical ozone maximum, which is instead best explained by a new mechanism: the *regime transition theory* (Section 4). In the regime transition theory, peak  $[O_3]$  occurs at an altitude around 26 km precisely because this marks the transition from an O-damped regime aloft, within which ozone is increasing towards the surface, to an  $O_3$ -damped regime below, within which ozone is decreasing towards the surface. We present an analytical expression for an idealized ozone profile under gray radiative transfer that produces an interior maximum of ozone at a self-consistent regime transition, and which improves intuition for the response of the Chapman+2 model to changes in UV (Section 6).

## 2 The Chapman+2 model with destruction by catalytic cycles and transport

A critical evaluation of the ozone maximum requires a model that can quantify the structural effects of ozone sinks arising from the Chapman cycle, catalytic cycles, and transport. We briefly introduce the Chapman cycle, which we then augment with two sinks representing catalytic cycles and transport. The Chapman cycle reactions are:



Reactions R2 and R4 depend on collisions, where M is a third body whose number density is that of air ( $n_a$ ). The collisional reactions proceed as the number density of the chemical reactants multiplied by a rate coefficient  $k_i$ ,  $i=2,4$ , e.g., reaction 2 has a rate of  $k_2[\text{O}][\text{O}_2][\text{M}]$ , which in general depends on temperature. Reactions R1 and R3 are photolysis reactions, and proceed as number density of the photolyzed species multiplied by the photolysis rate ( $J_{\text{O}_2}$  or  $J_{\text{O}_3}$ ). Photolysis rates couple chemistry and radiation together as follows:

$$J_{\text{O}_2}(z) = \int_{\lambda} q_{\text{O}_2}(\lambda) \sigma_{\text{O}_2}(\lambda) I(z, \lambda) d\lambda \quad (1)$$

$$J_{\text{O}_3}(z) = \int_{\lambda} q_{\text{O}_3}(\lambda) \sigma_{\text{O}_3}(\lambda) I(z, \lambda) d\lambda \quad (2)$$

with wavelength  $\lambda$ , quantum yield  $q_i(\lambda)$  (molecules decomposed per photon absorbed by species  $i$ ), absorption coefficient  $\sigma_i(\lambda)$  ( $\text{cm}^2 \text{ molec}^{-1}$ ) (shown in Fig. 3b), and UV flux density with respect to wavelength  $I(z, \lambda)$  ( $\text{photons cm}^{-2} \text{ s}^{-1} \text{ nm}^{-1}$ ). Top-of atmosphere UV flux ( $I(\infty, \lambda)$ ) is shown in Fig. 3a. Photolysis attenuates the UV flux:

$$I(z, \lambda) = I(\infty, \lambda) \exp\left(-\frac{\tau(z, \lambda)}{\cos \theta}\right) \quad (3)$$

where  $\tau(z, \lambda)$  is the optical depth as a function of wavelength, and  $\theta$  is the solar zenith angle, hereafter taken to be overhead sun for simplicity, so  $\cos \theta = 1$ . Because both  $\text{O}_2$  and  $\text{O}_3$  absorb UV, the optical depth at a given altitude depends on column-integrated  $\text{O}_2$  and  $\text{O}_3$  above that level:

$$110 \quad \tau(z, \lambda) = \sigma_{\text{O}_2}(\lambda) \chi_{\text{O}_2}(z) + \sigma_{\text{O}_3}(\lambda) \chi_{\text{O}_3}(z) \quad (4)$$

with  $\chi_{\text{O}_2}(z)$  giving the overhead column  $\text{O}_2$  ( $\int_z^{\infty} [\text{O}_2] dz$ ), and  $\chi_{\text{O}_3}(z)$  giving the overhead column  $\text{O}_3$  ( $\int_z^{\infty} [\text{O}_3] dz$ ).

## 2.1 The Chapman+2 model reactions

The Chapman cycle neglects the dominant sinks of ozone from catalytic chemistry and transport (Bates and Nicolet, 1950; Crutzen, 1970; Jacob, 1999; Brasseur and Solomon, 2005). These sinks involve photochemical reactions and transport among a system of at least tens of significant constituents. The consequences of the additional sinks of ozone from these processes is that ozone is approximately halved compared to in the Chapman cycle. Thus, calculating accurate photolysis rates, which depend on overhead column ozone, requires an accurate representation of basic state ozone as it is affected by these sinks. But, while the effects of these sinks are essential, many of their details are not thought to be part of a minimum essential explanation for the interior maximum of ozone. Therefore, we will parameterize the effects of these sinks on O and O<sub>3</sub>, facilitating a simple and tractable theory with a realistic basic state ozone profile. These sinks are parameterized by augmenting the Chapman cycle with two linear damping reactions that destroy O and O<sub>3</sub> respectively:



Representing these sinks as a linear damping is equivalent to adding an extra sink of O and O<sub>3</sub> in the form of a first-order decomposition reaction (analogous to radioactive decay). These sinks represent two pathways for the destruction of odd oxygen: destruction of odd oxygen can scale with atomic oxygen, as in R5 that proceeds at the rate  $\kappa_{\text{O}}[\text{O}]$ , or it can scale with ozone, as in R6 that proceeds at the rate  $\kappa_{\text{O}_3}[\text{O}_3]$ .

These reactions can be incorporated into the Chapman cycle to yield a Chapman+2 model of tropical stratospheric ozone, with the following prognostic equations for O and O<sub>3</sub>:

$$130 \quad \frac{\partial[\text{O}]}{\partial t} = 2J_{\text{O}_2}[\text{O}_2] - k_2[\text{O}][\text{O}_2][\text{M}] + J_{\text{O}_3}[\text{O}_3] - k_4[\text{O}][\text{O}_3] - \kappa_{\text{O}}[\text{O}] \quad (5)$$

$$\frac{\partial[\text{O}_3]}{\partial t} = k_2[\text{O}][\text{O}_2][\text{M}] - J_{\text{O}_3}[\text{O}_3] - k_4[\text{O}][\text{O}_3] - \kappa_{\text{O}_3}[\text{O}_3] \quad (6)$$

When solving for the ozone profile in the Chapman+2 model, there is generally several orders of magnitude more O<sub>2</sub> than odd oxygen ( $O_x \equiv \text{O} + \text{O}_3$ ), so for simplicity, O<sub>2</sub> will be treated as external to the Chapman+2 model, with fixed molar fraction of  $C_{\text{O}_2} = 0.21$ . Under this assumption, it is possible to solve for [O] and [O<sub>3</sub>] in steady state by setting  $\partial[\text{O}]/\partial t = \partial[\text{O}_3]/\partial t = 0$  in Equations 5 and 6, and solving for this system of two equations in two variables (O and O<sub>3</sub>):

$$[\text{O}_3] = \frac{J_{\text{O}_2} k_2}{k_4} C_{\text{O}_2}^2 n_a^3 \frac{1}{J_{\text{O}_3}[\text{O}_3] + J_{\text{O}_2} C_{\text{O}_2} n_a + \frac{\kappa_{\text{O}_3}[\text{O}_3]}{2} + \frac{J_{\text{O}_3} \kappa_{\text{O}}}{2k_4} + \frac{k_2 \kappa_{\text{O}_3} C_{\text{O}_2} n_a^2}{2k_4} + \frac{\kappa_{\text{O}} \kappa_{\text{O}_3}}{2k_4}} \quad (7)$$

where square brackets indicate number density (molec cm<sup>-3</sup>), and  $n_a$  is the number density of air. This equation is quadratic in  $[O_3]$  and mathematically implicit in height due to the dependence of  $J_{O_2}$  and  $J_{O_3}$  on ozone aloft. Note that  $J_{O_3}$  appears in the denominator as a photolytic sink of ozone.

140 An accompanying diagnostic equation for atomic oxygen is as follows:

$$[O] = \frac{J_{O_2} C_{O_2} n_a + J_{O_3} [O_3] + \frac{\kappa_{O_3} [O_3]}{2}}{k_2 C_{O_2} n_a^2 + \frac{\kappa_O}{2}} \quad (8)$$


In the absence of catalytic cycles and transport, i.e.,  $\kappa_O = \kappa_{O_3} = 0$ , Eqs. 7 and 8 reduce to the Chapman cycle (e.g., as analyzed in Craig, 1965). Yet, because the Chapman cycle is known to overestimate  $O_3$  by approximately a factor of two, these damping rates will be crucial for correcting these biases and leading to a realistic basic state. These damping rates will  
 145 be constrained based on prior knowledge of catalytic cycles and transport.

## 2.2 Constraining the Chapman+2 damping rates

Neither transport nor catalytic cycles generally act as a linear damping in all parts of the atmosphere or in all photochemical regimes. However, we will argue that the tropical stratosphere is in a regime where they can be fruitfully parameterized as such, facilitating theoretical insight.

150 Transport does not generally act as a linear damping, and indeed the Brewer-Dobson circulation is known to be a source of ozone in the extratropics (e.g., Dobson, 1956). However, in the tropical lower stratosphere, where transport might in principle be represented as a leaky tropical pipe (Neu and Plumb, 1999) such as in Match and Gerber (2022), its effects can be approximated as a linear damping in order to understand peak  $[O_3]$ . This linear damping results because ozone is being constantly upwelled from an ozone-poor region (the tropical tropopause layer) into a region over which it decays with a characteristic  
 155 scale height (Brasseur and Solomon, 2005, Section 3.5.2). And, because transport is only important for ozone in the tropical lower stratosphere and not farther aloft (e.g., Garcia and Solomon, 1985; Perliski et al., 1989), a fact that will emerge self-consistently within the Chapman+2 model, parameterizing the effects of transport as a constant damping throughout the tropical stratosphere can lead to an accurate representation in the tropical lower stratosphere without imposing significant errors farther aloft. We consider that transport leads to a relaxation rate that scales with  $\bar{w}^* = 0.3 \text{ mm s}^{-1}$  divided by a reference  
 160 vertical scale of approximately 2 km, leading to a damping rate of  $\kappa_{\bar{w}^*} = (3 \text{ months})^{-1}$ . For consistency, this damping will be applied to O and  $O_3$ , although it will be found to only significantly affect  $O_3$  given the short photochemical lifetime of O.

Like transport, catalytic cycles also do not generally act as a linear damping. This is because they involve two- (and sometimes three-)body reactions whose rates depend on the abundance of the catalysts, which are often co-evolving with the overall photochemical state. Thus, in order to treat catalytic destruction of ozone as a linear damping with a steady, altitude-dependent  
 165 damping rate, we assume that the number density of the catalysts and the temperature-dependent reaction rates are constant. We then must use these constant profiles of damping rates to damp O and  $O_3$ . In order to determine these damping rate profiles, and whether they damp O or  $O_3$ , we distinguish catalytic cycles driven by some catalyst Z by their net effects, where representative cycles leading to each net effect are shown below:



170 The most significant catalysts driving each class of catalytic cycle are as follows (e.g., Brasseur and Solomon, 2005): destruction of  $O_3$  is driven by  $Z = OH$ , destruction of  $O$  is driven by  $Z = H$ , and destruction of  $O + O_3$  is driven by  $Z = H, OH, NO, Cl,$  and  $Br$ . In the tropical stratosphere, these latter reactions that destroy  $O + O_3$  tend to be rate-limited by the destruction of  $O$ . This rate-limitation arises because reaction  $Z + O_3$  produces  $ZO$ , which is often photolyzed to complete a null cycle, so only if  $ZO$  reacts with  $O$  does the catalytic cycle ultimately destroy two  $O_x$ . Therefore, such cycles are considered to damp odd  
175 oxygen at the rate of  $2k_{ZO+O}[ZO]$ . Catalytic reaction rates are taken from Brasseur and Solomon (2005).

Combining the effects of transport and catalytic cycles leads to these estimates for the damping rates:

$$\kappa_O = \kappa_{\bar{w}^*} + a_5[OH] + a_7[HO_2] + 2b_3[NO_2] + 2d_3[ClO] + 2e_3[BrO] \quad (9)$$

$$\kappa_{O_3} = \kappa_{\bar{w}^*} + a_2[H] + a_6[OH] + a_{6b}[HO_2] \quad (10)$$

where we have referred to the reaction rate coefficients ( $a_5, a_7, b_3,$  etc.) as in Brasseur and Solomon (2005).

180 As a link to existing frameworks, the catalytic component of these damping rates can be related to the budget of generalized odd oxygen ( $O_y$ ), which was defined in Brasseur and Solomon (2005) to include a broader set of chemical constituents that can serve as reservoirs for odd oxygen under stratospheric photochemistry<sup>2</sup>. Equations 9 and 10 include all of the dominant sinks of  $O_y$  that are linear in  $O$  or  $O_3$ . These damping rates treat the concentrations of catalysts as constant and neglect conversions of generalized odd oxygen between reservoir species, so do not provide an exhaustive account of the  $O_y$  budget. Nonetheless,  
185 they will serve to effectively parameterize the sinks of  $O$  and  $O_3$ .

Profiles for these damping rates  $\kappa_O$  and  $\kappa_{O_3}$  are estimated by using globally-averaged vertical profiles for the chemical constituents from the chemistry-climate model SOCRATES (Brasseur et al., 1990), as tabulated in Brasseur and Solomon (2005). The damping rates are approximated crudely insofar as catalyst profiles in the tropics are approximated with their

<sup>2</sup>On their page 414,  $O_y$  is defined as  $O(^3P) + O(^1D) + O_3 + NO_2 + 2NO_3 + HNO_3 + HO_2NO_2 + 2N_2O_5 + ClO + 2Cl_2O_2 + 2OCIO + 2ClONO_2 + BrO + 2BrONO_2$ .



global average. The resulting profiles of damping rates are shown in Fig. 3c. These damping rate profiles can be validated  
190 against an independent estimate of the photochemical damping timescale from the chemical transport model MOBIDIC (as  
calculated for the Cariolle v2.9 linear ozone model, personal communication with Cariolle, 2023, dashed cyan curve in Fig.  
3c). These linear ozone model coefficients are closely related to an effective damping rate of odd oxygen excluding the effects  
of transport, analogous in our framework to the quantity  $\kappa_{O_x, \text{eff}} = (\kappa_{O_3} - \kappa_{\bar{w}^*}) + [O]/[O_3] * (\kappa_O - \kappa_{\bar{w}^*})$  (solid cyan curve in Fig.  
3c). These two cyan curves of the photochemical damping timescale are approximately consistent in magnitude and vertical  
195 structure, building confidence in our damping rates.

### 2.3 Evaluating O<sub>3</sub> in the Chapman+2 model

Prescribing these damping rates  $\kappa_O$  and  $\kappa_{O_3}$ , it is then possible to solve Equations 7 and 8 by integrating from the top of  
the atmosphere downwards, solving jointly for O<sub>3</sub> and the UV fluxes at each vertical level. Further details of our numerical  
approach are in Appendix A.


200 Fig. 3 compares numerical solutions of the Chapman cycle and Chapman+2 model compared to observed tropical [O<sub>3</sub>]. As  
is well known, the Chapman cycle overestimates ozone by approximately a factor two (Fig. 3d, gray vs. black). These overes-  
timates are significantly mitigated in the Chapman+2 model (Fig. 3d, magenta vs. black). The improved ozone magnitudes in  
the Chapman+2 model allow UV flux to penetrate more deeply than in the Chapman cycle (Fig. 3e,f), leading to more realistic  
photolysis rates.

205 Agreement between the Chapman+2 model and observations is imperfect, which is unsurprising given that this work employs  
many simplifying approximations. Many of these approximations are required to subsequently derive an explicit analytical  
expression to the Chapman+2 model ozone profile. For example, we will present results for overhead sun impinging on an  
isothermal atmosphere, although our model can also be run at other solar zenith angles or with vertically-varying temperature.  
We have also approximated transport and catalytic cycles as a linear damping, used globally-averaged catalytic profiles, and  
210 neglected optical scattering. Despite these approximations, the Chapman+2 model produces a reasonable fit to the observed  
profile, and will be considered to produce a credible interior maximum of ozone. The remainder of the paper seeks to explain  
why the Chapman+2 model produces an interior maximum.

## 3 Understanding the ozone maximum

Understanding how the Chapman+2 model produces an interior maximum is challenging when considering the ozone number  
215 density in Eq. 7, so we perform a scale analysis to identify the dominant photochemical-transport regimes at different altitudes.  
Three limits can be defined based on whether the sink of odd oxygen is dominated by the Chapman cycle sink from the  
reaction O + O<sub>3</sub>, the damping of O, or the damping of O<sub>3</sub>. These limits correspond to different dominant terms in the six-term  
denominator of Eq. 7.

If the Chapman cycle sink of ozone dominates, then the dominant term in the denominator of Eq. 7 is  $J_{O_3}[O_3]$ , and ozone  
220 scales as:



**Figure 3.** Boundary conditions and solutions to the Chapman cycle and Chapman+2 model for tropical ozone. (a) UV flux at the top of the atmosphere. (b) Absorption coefficients for O<sub>2</sub> and O<sub>3</sub>. (c) Damping rates of O (red) and O<sub>3</sub> (blue) estimated from Eqs. 9 and 10, using catalyst profiles from the chemistry-climate model SOCRATES, as tabulated in Brasseur and Solomon (2005). The effective damping rate of O<sub>x</sub> (solid cyan) is comparable to the derived O<sub>3</sub> relaxation rate in the chemical transport MOBIDIC as calculated in the Cariolle v2.9 linear ozone model (dashed cyan). (d) Ozone profile in numerical solutions to the Chapman cycle. Numerical solutions are compared to satellite-observed ozone from the SWOOSH dataset averaged from 1984-2023 and from 30°S-30°N (black). (e) UV flux for the Chapman+2 experiment and (f) Chapman cycle experiment ( $\kappa_{\text{O}} = \kappa_{\text{O}_3} = 0$ , where we indicate the level of unit optical depth ( $\tau(\lambda) = 1$ ) in gray). For clarity, wavelength axes are restricted to 180-320 nm although the numerical solution extends to 800 nm into the weakly-absorbing Chappuis bands. A similar introductory figure appears in Match et al. (2024).

$$[\text{O}_3] = \left( \frac{J_{\text{O}_2} k_2}{J_{\text{O}_3} k_4} \right)^{1/2} C_{\text{O}_2} n_a^{3/2} \quad (11)$$

Eq. 11 reproduces the well-known Chapman cycle limit, where the dominant reactions are R1-R4, as presented in most textbook explanations for the shape of the ozone layer. The vertical structure of the ozone layer in Eq. 11 arises predominantly from the number density of air,  $n_a^{3/2}$  (assumed invariant under Chapman photochemistry), and from the ratio of photolysis rates  
 225  $(J_{O_2}/J_{O_3})^{1/2}$ . The presence of  $J_{O_3}$  in the denominator indicates that photolysis of  $O_3$  is an effective sink of  $O_3$  by producing atomic oxygen that can then destroy  $O_3$  through R4. We refer to this as a *photolytic sink*. The fact that photolysis of  $O_3$  acts as a photolytic sink might seem surprising since it is typically understood to not affect ozone due to the strong null cycle  $R2 \rightleftharpoons R3$ . However, that null cycle has some leakage into R4. Thus, even though most of the photolysis of ozone does not lead to the destruction of ozone (legitimizing the concept of odd oxygen), most of the destruction of ozone requires photolysis of ozone,  
 230 in order to supply atomic oxygen—hence  $J_{O_3}$  suppresses ozone as a photolytic sink.

If the damping of O dominates through  $J_{O_3}\kappa_O/2k_4$ , ozone number density scales as:

$$[O_3]_{O\text{-damped}} = \frac{2J_{O_2}k_2C_{O_2}^2n_a^3}{J_{O_3}\kappa_O} \quad (12)$$


Equation 12 corresponds to the limit where the dominant reactions are R1-R3 and R5. The Chapman cycle and O-damped regime (Eqs. 11 and 12) share key structural aspects, as ozone scales as  $((J_{O_2}/J_{O_3})n_a^3)^n$ , where  $n = 1/2$  in the Chapman cycle  
 235 regime, and  $n = 1$  in the O-damped regime. Note that in both cases, photolysis of  $O_3$  appears in the denominator as a photolytic sink that is necessary for producing atomic oxygen that can either react with ozone (in the Chapman cycle) or be damped (in the O-damped regime). Thus, these regimes both derive their structure from a photolytic sink, and both satisfy the source/sink competition paradigm.

If the damping of  $O_3$  dominates through  $k_2\kappa_{O_3}C_{O_2}n_a^2/2k_4$ , ozone number density scales as:

$$240 \quad [O_3]_{O_3\text{-damped}} = \frac{2J_{O_2}[O_2]}{\kappa_{O_3}} \quad (13)$$

Equation 13 corresponds to the limit where the dominant reactions are R1-R3 and R6. In the  $O_3$ -damped regime, ozone destruction does not depend on photolysis of ozone, which therefore does not appear in the ozone equation. With this sink that is independent of photolysis, ozone scales with the production rate divided by the damping rate of  $O_3$ , consistent with the source-controlled paradigm. Note that, in this regime, R3 can still be fast compared to the production of odd oxygen from R1,  
 245 but it has negligible effect on ozone concentrations because, to leading order, it drives the null cycle  $R2 \rightleftharpoons R3$ , i.e., not only does most photolysis of ozone not lead to destruction of ozone, as is generally the case, but also most destruction of ozone does not involve photolysis of ozone, which is not true in the Chapman cycle or O-damped regime.

Thus, the prevailing textbook paradigms for explaining the interior maximum of ozone correspond to well-defined limits of the Chapman+2 model corresponding to either the Chapman cycle regime or O-damped regime (source/sink competition  
 250 paradigm, Fig. 1b) or to an  $O_3$ -damped regime (source-controlled paradigm, Fig. 1a). Which regime actually prevails is an empirical question. Fig. 3c reveals that the damping of O is everywhere larger than the damping of  $O_3$ , but this does not imply that the ozone layer is everywhere in an O-damped regime because the altitude-dependent partitioning between [O] and [ $O_3$ ]



**Figure 4.** (a) Odd oxygen sink regimes due to catalytic chemistry and transport based on the effective damping rates of O and O<sub>3</sub> in the tropics. Above 26 km, the ozone layer is in an O-damped regime (red region). Below 26 km, the ozone layer is in an O<sub>3</sub> damped regime (blue region). The Chapman cycle sink does not dominate anywhere. (b) Fraction of  $\kappa_O$  contributed by each component in Eq. 9. (c) Fraction of  $\kappa_{O_3}$  contributed by each component in Eq. 10.

also matters. Determining the O-damped versus O<sub>3</sub>-damped regimes requires evaluating the dominant terms in the denominator of the catalytic ozone solution (Eq. 7), where the contribution of the Chapman cycle sink scales as  $J_{O_3}[O_3]$ , the damping of O scales as  $J_{O_3}\kappa_O/2k_4$ , and damping of O<sub>3</sub> scales as  $k_2\kappa_{O_3}C_{O_2}n_a^2/2k_4$ .

The vertical profile of catalytic regimes can be categorized as follows (Figure 4a):

- Above 26 km, the stratosphere is dominated by O-damping. If the interior maximum of ozone occurred well above 26 km, it would be explained by the source/sink competition paradigm.
- Below 26 km, the stratosphere is dominated by O<sub>3</sub>-damping. If the interior maximum of ozone occurred well below 26 km, it would be explained by the source-controlled paradigm.

Yet, the interior maximum of ozone in the Chapman+2 model occurs exactly at this transition, at an altitude of 26 km, hinting at the need to consider both regimes.

The damping of O and O<sub>3</sub> that establishes each regime can be further decomposed into additive contributions from the terms in Eqs. 9 and 10 (Figs. 4b and 4c). Throughout the stratosphere of the Chapman+2 model, the damping of O is dominated by NO<sub>2</sub> (Fig. 4b, red curve). Towards the stratopause at 50 km, the stratosphere is in an O-damped regime, but our model overestimates the NO<sub>x</sub> sink and therefore fails to correctly identify that the HO<sub>x</sub> sink should dominate at these altitudes (as in,

e.g., Brasseur and Solomon, 2005). Lower in the stratosphere, near peak  $[O_3]$ , the Chapman+2 model correctly captures the dominance of the  $NO_x$  sink, where the damping rate of O can be accurately approximated as  $\kappa_O \approx b_3[NO_2]$ . The damping of  $O_3$  is dominated by H in the upper stratosphere, by OH lower down around 40 km, and by transport below 35 km. The dominance of transport in the lower stratosphere means that, in the  $O_3$ -damped regime below 26 km,  $\kappa_{O_3} \approx \kappa_{\bar{w}}$  (Fig. 4c, black curve). Thus, the odd oxygen sink regimes can be interpreted as transitioning from an O-damped regime dominated by  $NO_x$  above 26 km to an  $O_3$ -damped regime dominated by transport below 26 km.

#### 4 A new theory for the tropical ozone maximum


In the Introduction, we showed that the interior maximum of observed tropical  $[O_3]$  could not be reproduced by the scaling relationships from either the source-controlled paradigm or the source/sink competition paradigm. Now equipped with the Chapman+2 model and scaling relationships for ozone in each sink regime, we reaffirm that the interior maximum cannot be explained by either paradigm.

Fig. 5 shows the observed  $[O_3]$  profile (black) compared to ozone in the Chapman+2 model (magenta) and its limits in the O-damped regime (solid red) and  $O_3$ -damped regime (solid blue). Above 26 km,  $[O_3]$  closely follows the scaling from the O-damped regime (solid red). Below 26 km,  $[O_3]$  closely follows the scaling from the  $O_3$ -damped regime (solid blue). To examine where each regime predicts peak  $[O_3]$ , these theoretical scalings can be artificially extended beyond where they formally apply (dashed curves). When the O-damped regime is extended downwards (red dashed), it predicts an interior maximum at 15 km, far below the ozone maximum and its range of applicability. When the  $O_3$ -damped regime is extended upwards (blue dashed), it predicts an interior maximum at 35 km, far above the ozone maximum and its range of applicability. These predictions for peak  $[O_3]$  are similar to those shown in Figure 2, except now they are formulated as quantitative limits of the Chapman+2 model, they use self-consistent photolysis rates as part of the Chapman+2 model solution, and they include (modest) altitude-dependent contributions from  $\kappa_O$  and  $\kappa_{O_3}$ . Figure 2 first suggested limitations of the prevailing paradigms, and Figure 5 confirms that these limitations verge on being paradoxical: each textbook paradigm is capable of producing an interior maximum, but these maxima occur at the wrong altitude and in a region where they do not apply.

We propose a new theory for the interior maximum: tropical  $[O_3]$  peaks around 26 km because this marks the transition from an O-damped regime aloft, within which  $[O_3]$  is increasing towards the surface, to an  $O_3$ -damped regime below, within which  $[O_3]$  is decreasing towards the surface. The *regime transition theory* is illustrated in Fig. 6.

#### 5 Why there is a regime transition

Although it has not previously been invoked to explain peak  $[O_3]$ , it is well known that there is a regime transition in ozone photochemistry between 25 and 30 km from a photochemically-dominated regime aloft to a transport-dominated regime below (Garcia and Solomon, 1985; Perliski et al., 1989; Brasseur and Jacob, 2017). This regime transition has been fruitfully interpreted in terms of the equilibration timescale for odd oxygen. Aloft, odd oxygen equilibrates with respect to photochemistry




**Figure 5.** The Chapman+2 model (magenta) compared to observed tropical  $O_3$  from SWOOSH averaged from 1984-2023 and from 30°S-30°N (black). Above 26 km, ozone follows the theoretical scaling for the O-damped regime (solid red curve). From 26 km down to the troposphere, ozone follows the theoretical scaling of the  $O_3$ -damped regime (solid blue curve). Extending the theoretical scalings across the whole domain (dashed curves) reveals the apparent paradox that each scaling predicts ozone to maximize outside its region of applicability. This is resolved by the regime transition theory: ozone maximizes at the transition from an O-damped regime aloft to an  $O_3$ -damped regime below.

very rapidly, quickly erasing any anomalies induced by the transport of odd oxygen. This equilibration timescale becomes more sluggish towards lower altitudes, where transport is then able to generate anomalies from photochemical equilibrium.

300 This transition is closely analogous to the transition considered herein from an O-damped regime to an  $O_3$ -damped regime. In order to understand either regime transition, the question becomes: why does the photochemical timescale become longer than the transport timescale below some altitude, or relatedly, why does the O-damping become weaker than the  $O_3$ -damping?

This question can be answered by assessing the contributions to the structure of O-damping versus of  $O_3$ -damping. Using the terms from the denominator of Equation 7, we define  $\gamma_O$  as a non-dimensional ratio measuring the strength of O-damping

305 compared to  $O_3$ -damping, defined as follows:




**Figure 6.** The source-controlled paradigm (right) predicts peak  $[O_3]$  too high and where it is inapplicable. The source/sink competition paradigm predicts peak  $[O_3]$  too low and where it is inapplicable. The regime transition theory (center) explains that  $[O_3]$  peaks at the transition between these two regimes.

$$\gamma_0 = \frac{J_{O_3} \kappa_O}{k_2 \kappa_{O_3} C_{O_2} n_a^2} \quad (14)$$

Under realistic conditions in which the ozone layer is either O-damped or  $O_3$ -damped, i.e., the Chapman cycle sink does not dominate, then when  $\gamma_0 > 1$ , the ozone layer is O-damped, and when  $\gamma_0 < 1$ , the ozone layer is  $O_3$ -damped. The magnitude of  $\gamma_0$  generally declines from the upper stratosphere downward, and where it crosses below 1 is, by construction, the regime transition.

The terms that contribute most to the vertical structure of  $\gamma_0$  are plotted in Figure 7. The dominant driver of the regime transition is the rise in air density towards lower altitudes. Larger air density contributes to the regime transition by speeding up R2, thereby partitioning odd oxygen away from O and in favor of  $O_3$ . Partitioning odd oxygen away from O reduces the magnitude of O-damping (which scales as  $\kappa_O[O]$ ) and strengthens  $O_3$ -damping (which scales as  $\kappa_{O_3}[O_3]$ ). The effects of the rising air density are quadratic because this partitioning scales with  $[O_2]$  and  $[M]$ , both of which are proportional to air density. There are also two smaller, but still significant, drivers of the regime transition. The decline in the photolysis rate of  $O_3$  towards lower altitudes helps drive the regime transition by repartitioning odd oxygen towards  $O_3$  and away from O, favoring  $O_3$ -damping at the expense of O-damping. The decline in  $\kappa_O$  from 35 km down to 26 km also favors the regime transition, reflecting that  $[NO_2]$  peaks at 35 km.

The decline in  $\kappa_{O_3}$  towards lower altitudes modestly opposes the regime transition. Because we have assumed that the damping from transport is uniform, this decline must result from catalytic sinks, primarily the drop-off in  $[OH]$  and  $[H]$ .



**Figure 7.** Multiplicative factors contributing to the vertical structure of  $\gamma_O$  in the tropics (defined in text, plotted in black), relative to where  $\gamma_O$  reaches unity at 26 km and therefore defines the regime transition. Factors that decline from above 26 km to below 26 km drive the regime transition, so the dominant driver is air density (cyan). Smaller contributions come from the photolysis rates of  $O_3$  (magenta) and the damping rate of  $O$  (red). Modestly opposing the regime transition is the damping rate of  $O_3$  (blue).

Thus, there is a transition from an  $O$ -damped regime aloft to an  $O_3$ -damped regime below primarily because, descending towards the ozone maximum, the atmosphere tends to repartition odd oxygen away from  $O$  and towards  $O_3$ . This repartitioning inhibits damping of  $O$  and invigorates damping of  $O_3$ , which eventually dominates. This repartitioning occurs primarily due to  
 325 increasing air density, but also due to attenuation of UV that photolyzes  $O_3$ .

## 6 An explicit solution for the ozone layer under gray radiation

The regime transition theory suggests a two-regime conceptual model for tropical stratospheric ozone: ozone increases towards its interior maximum in an  $O$ -damped regime before reaching a transition altitude at which it maximizes, below which it decreases towards the tropopause in an  $O_3$ -damped regime. Conceiving of the ozone layer in terms of an  $O$ -damped regime,  
 330 a transition altitude, and an  $O_3$  damped regime can improve conceptual understanding of its basic state and its sensitivity to perturbations. In this section, we focus on the basic state by showing that this conceptual understanding can be encoded into an explicit analytical expression for the ozone profile that produces an interior maximum at the regime transition. Then, in the following section, we will consider the sensitivity to perturbations.

We preface our derivation of an explicit expression for the profile of ozone under idealized boundary conditions by first  
 335 noting that there are no previously published mathematically explicit expressions for the ozone profile under any set of as-



sumptions, let alone those that would produce an interior maximum at a regime transition. This is due to two key obstacles: (1) ozone photochemistry is mathematically implicit, and (2) it relies on spectral integrals across non-analytical functions. The obstacle arising from spectral integrals across non-analytical functions is generic to radiative transfer problems. Yet, recent work has advanced understanding of the emergent effects of longwave radiative transfer by judiciously approximating non-analytical absorption spectra for CO<sub>2</sub> or H<sub>2</sub>O with analytical functions, leading to *simple spectral models* (SSMs, after Jeevanjee and Fueglistaler, 2020) that can then be coupled to other aspects of climate dynamics (Jeevanjee and Fueglistaler, 2020; Jeevanjee et al., 2021; Pierrehumbert, 2011; Romps et al., 2022). Here, we develop simple spectral models for ozone photochemistry, in certain limits of which the UV radiative transfer becomes mathematically explicit. Under these and other approximations, mathematically explicit ozone profiles can be derived under the regime transition theory (this section), as well as for the source/sink competition paradigm (Appendix B) and the source-controlled paradigm (Appendix C)

In each derivation, we begin by assuming gray radiation with uniform and fully-overlapping absorption by O<sub>2</sub> and O<sub>3</sub> across a spectral window  $\mu$  (nm) with absorption coefficients  $\sigma_{\text{O}_2}^*$  and  $\sigma_{\text{O}_3}^*$ . We also assume overhead sun, an isothermal atmosphere, uniform  $\kappa_{\text{O}}$  and  $\kappa_{\text{O}_3}$ , and unit quantum yields ( $q_{\text{O}_2}^* = q_{\text{O}_3}^* = 1$ ).

### 6.1 Upper branch in an O-damped regime

To solve for an ozone profile under the regime transition theory, we must solve from the top down, beginning in the O-damped regime. Under gray radiation, the photolysis rates can be expressed implicitly as a function of ozone by substituting into Eqs. 1 and 2:

$$J_{\text{O}_2}(z) = \mu \sigma_{\text{O}_2}^* I_{\infty} q_{\text{O}_2}^* \exp(-\sigma_{\text{O}_2}^* \chi_{\text{O}_2}(z) - \sigma_{\text{O}_3}^* \chi_{\text{O}_3}(z)) \quad (15)$$

$$J_{\text{O}_3}(z) = \mu \sigma_{\text{O}_3}^* I_{\infty} q_{\text{O}_3}^* \exp(-\sigma_{\text{O}_2}^* \chi_{\text{O}_2}(z) - \sigma_{\text{O}_3}^* \chi_{\text{O}_3}(z)) \quad (16)$$

The photolysis rates depend on column ozone, so it would seem that the ozone profile should depend implicitly on ozone aloft. However, the O-damped regime is in a photolytic sink regime where ozone scales with the ratio  $J_{\text{O}_2}/J_{\text{O}_3}$ , which under gray radiation reduces to the ratio of the absorption coefficients  $\sigma_{\text{O}_2}^*/\sigma_{\text{O}_3}^*$ . This leads to an explicit solution for ozone in the O-damped regime (Eq. 12):

$$[\text{O}_3]_{\text{gray,O-damped}} = \frac{2\sigma_{\text{O}_2}^* k_2 C_{\text{O}_2}^2 n_a^3}{\sigma_{\text{O}_3}^* \kappa_{\text{O}}} \quad (17)$$

The only altitude-dependence is that  $[\text{O}_3]$  scales with  $n_a^3$ . Thus, absent a transition to an O<sub>3</sub>-damped regime, the O-damped ozone layer would increase all the way down and have no interior maximum. Equation 17 can be integrated to yield column ozone:

$$\chi_{O_3}(z)|_{\text{gray,O-damped}} = \frac{H}{3} [\text{O}_3]_{\text{gray,O-damped}} \quad (18)$$

This expression for the column ozone under gray radiation and O-damping can then be substituted back into the photolysis rates to solve explicitly for  $J_{O_2}$  (Eq. 15) and  $J_{O_3}$  (Eq. 16).

This derivation reveals a more general result, which is that  $[\text{O}_3]$  does not have an interior maximum within a photolytic sink regime under gray radiation. This applies to both the O-damped regime and the Chapman cycle. In order to be correct, explanations for the interior maximum of  $[\text{O}_3]$  within the Chapman cycle must invoke spectral structure of absorption coefficients. To see how the spectral structure of absorption coefficients can lead to an interior maximum of ozone within the Chapman cycle or the O-damped limit, we present a two-band simple spectral model that supports an analytical expression for an interior maximum of ozone in the source/sink competition paradigm in Appendix B.

## 6.2 Regime transition and peak $[\text{O}_3]$

The O-damped regime continues downwards until reaching a regime transition, which must be self-consistently calculated to occur where the damping of O and  $\text{O}_3$  are exactly co-dominant, i.e., when the non-dimensional parameter  $\gamma_O$  defined in Eq. 14 equals unity. For Earth-like parameters, the regime transition also corresponds to the peak  $[\text{O}_3]$ . To solve analytically for the regime transition, it must be assumed (realistically) that the dominant absorber of UV is ozone. Under that assumption and using the column ozone scaling for the O-damped regime, the ozone photolysis rate scales as follows:

$$J_{O_3}(z) = \sigma_{O_3}^* q_{O_3}^* I_\infty \Delta\lambda \exp(-\sigma_{O_3}^* \chi_{O_3}(z)|_{\text{gray,O-damped}}) \quad (19)$$

Substituting this expression for the photolysis rate of ozone into the transition condition that  $\gamma_O = 1$  and solving for  $z$  yields the transition altitude:

$$z_t = H \left( \frac{1}{3} W \left( \frac{\tau_{O_2}(0) \alpha_O^{1/2}}{\alpha_{O_3}^{3/2}} \right) + \frac{1}{2} \ln \frac{\alpha_{O_3}}{\alpha_O} \right) \quad (20)$$

where  $W$  is the Lambert  $W$  function, which when evaluated at  $x$  returns the value  $w$  such that  $w \exp(w) = x$ , and we have defined the following three non-dimensional parameters of use for interpreting the transition altitude scaling:

$$\alpha_O \equiv \frac{\kappa_O}{k_2 C_{O_2} n_{a_0}^2} \quad (21)$$

$$\alpha_{O_3} \equiv \frac{\kappa_{O_3}}{\sigma_{O_3}^* q_{O_3}^* I_\infty \Delta\lambda} \quad (22)$$

$$\tau_{O_2}(0) = \sigma_{O_2}^* C_{O_2} n_{a_0} H \quad (23)$$

The first nondimensional parameter,  $\alpha_O$ , measures the strength of O-damping compared to the rate at which atomic oxygen combines with  $O_2$  to form  $O_3$  (R2). The second nondimensional parameter,  $\alpha_{O_3}$ , measures the strength of  $O_3$ -damping compared to the photolysis rate of  $O_3$  at the top of the atmosphere (R3). The third nondimensional parameter is the optical depth of  $O_2$  at the surface.

Substituting the expression for  $z_t$  into the scaling for ozone in the O-damped regime (Eq. 17) yields an analytical expression for ozone at the transition altitude:

$$[O_3](z_t) = \frac{2}{H\sigma_{O_3}^*} W \left( \frac{\alpha_O^{1/2} \tau_{O_2}(0)}{\alpha_{O_3}^{3/2}} \right) \quad (24)$$

This is an explicit analytical expression for  $O_3$  at the transition altitude, which for realistic parameters is also the peak  $[O_3]$ . Below the regime transition, the ozone profile shifts to the  $O_3$ -damped regime.

### 6.3 Lower branch in $O_3$ -damped regime

To solve for the lower branch of the  $O_3$  profile, we take advantage of the continuity of UV flux across the regime transition. Thus, our approach considers an  $O_3$ -damped region below  $z_t$  with constant  $\kappa_{O_3}$ .

In the  $O_3$ -damped regime, ozone scales with its production rate, and we solve for  $J_{O_2}$  by substituting the expression for  $O_3$  in the  $O_3$ -damped regime (Eq. 13) into the column ozone integral:

$$J_{O_2}(z) = \mu \sigma_{O_2}^* q_{O_2}^* I_\infty \exp(-\sigma_{O_2}^* \chi_{O_2}(z)) \exp \left( -\sigma_{O_3}^* \left( \chi_{O_3}(z_t) + \int_z^{z_t} \frac{2J_{O_2}[O_2]}{\kappa_{O_3}} dz \right) \right) \quad (25)$$

Taking the natural logarithm of both sides of Eq. 25 and differentiating with respect to  $z$  leads to a differential equation for  $J_{O_2}$  as a function of  $z$ :

$$\frac{dJ_{O_2}(z)}{dz} = \frac{2\sigma_{O_3}^* C_{O_2} n_{a_0}}{\kappa_{O_3}} J_{O_2}(z)^2 \exp(-z/H) + \sigma_{O_2}^* C_{O_2} n_{a_0} J_{O_2}(z) \exp(-z/H) \quad (26)$$

This first-order nonlinear ordinary differential equation can be solved by separation of variables and integrated from the transition altitude  $z_t$  downwards using the following boundary condition:

$$J_{O_2}(z_t) = \sigma_{O_2}^* \mu q_{O_2}^* I_\infty \exp(-\sigma_{O_2}^* \chi_{O_2}(z_t)) \exp(-\sigma_{O_3}^* \chi_{O_3}(z)) \Big|_{\text{gray, O-damped}} \quad (27)$$

which leads to an equation for  $J_{O_2}$ :

$$J_{O_2}(z) = \frac{\sigma_{O_2}^* \kappa_{O_3}}{2\sigma_{O_3}^* \left( \left( \frac{\sigma_{O_2}^* \kappa_{O_3}}{2J_{O_2}(z_t) \sigma_{O_3}^*} + 1 \right) \exp(\tau_{O_2}(0)(e^{-z/H} - e^{-z_t/H})) - 1 \right)} \quad (28)$$

410 This expression for  $J_{O_2}(z)$  can be substituted into the equation for  $O_3$  under  $O_3$ -damping to yield the ozone profile.

In Appendix C, we present an analytical solution to the ozone layer under gray radiation and strong damping such that  $z_t$  can be approximated as the top of the atmosphere. In the limit of large  $\kappa_{O_3}$ , this solution reproduces the “sweet spot” explanation for the ozone layer in the source-controlled paradigm.

## 6.4 Putting the pieces together

415 Summarizing, ozone in the upper branch is in an O-damped regime (Eq. 17) down to the altitude of the regime transition,  $z_t$  (Eq. 20), below which ozone is in an  $O_3$ -damped regime (inferred from Eq. 28). Putting the pieces of these regimes together yields an explicit analytical profile of ozone in the Chapman+2 model under gray radiation:

$$[O_3]_{\text{gray}} = \begin{cases} \frac{2\sigma_{O_2}^* k_2 C_{O_2}^2 n_{a_0}^3 \exp\left(\frac{-3z}{H}\right)}{\sigma_{O_3}^* \kappa_{O_3}} & \text{if } z \geq z_t \\ \frac{2}{H\sigma_{O_3}^*} W\left(\frac{\alpha_0^{1/2} \tau_{O_2}(0)}{\alpha_{O_3}^{3/2}}\right) & \text{if } z = z_t \\ \frac{\sigma_{O_2}^* C_{O_2} n_{a_0} \exp(-z/H)}{\sigma_{O_3}^* \left( \left( \frac{\sigma_{O_2}^* \kappa_{O_3}}{2J_{O_2}(z_t) \sigma_{O_3}^*} + 1 \right) \exp(\tau_{O_2}(0)(e^{-z/H} - e^{-z_t/H})) - 1 \right)} & \text{if } z < z_t \end{cases} \quad (29)$$


where we used the nondimensional parameters defined in Eqs. 21-23. The UV flux is continuous across the transition altitude, but ozone is not necessarily continuous across  $z_t$ . Note that the  $[O_3]$  at  $z_t$  is consistent between the O-damped regime (first line of Eq. 29) and the explicit solution at  $z_t$  (second line of Eq. 29).

Fig. 8 shows successive approximations to the ozone profile, beginning with the full Chapman+2 model solution and ending with the analytical solution with constant damping rates and gray radiation. The assumptions that move the system towards its analytical solution can be seen to degrade the solution at each step. However, even the analytical profile retains a realistic structure, with its details notably dependent on parameters selected both for their plausibility and their post hoc agreement with the observed profile. Rather than the details of the fit, the advantage of the gray solution is that it is the first simple spectral model of ozone that affords an explicit solution to an ozone profile that has an interior maximum. It provides a quantitative framework for considering the response to perturbations, as next considered briefly in the Discussion.

## 7 Discussion

### 430 7.1 Understanding the response to perturbations: the case of doubling UV

Distinguishing among competing theories for the same phenomenon can be justified, in part, if those theories make different predictions for the response to perturbations. This is the case among the two textbook paradigms and our new theory. This is



**Figure 8.** Beginning with observations compared to the full Chapman+2 model calculation with altitude-dependent  $\kappa_{\text{O}}$  and  $\kappa_{\text{O}_3}$  and spectrally-varying radiative transfer, we make successive approximations in our numerical solutions until reaching the analytical solution from Equation 29 with  $\kappa_{\text{O}} = 10^{-2} \text{ s}^{-1}$  and  $\kappa_{\text{O}_3} = (3 \text{ months})^{-1}$  and gray radiation with  $\sigma_{\text{O}_2} = 10^{-25} \text{ cm}^2 \text{ molec}^{-1}$  and  $\sigma_{\text{O}_3} = 5 * 10^{-18} \text{ cm}^2 \text{ molec}^{-1}$ .


illustrated by considering the ozone response to a spectrally uniform doubling of top-of-atmosphere UV flux (holding  $\kappa_{\text{O}}$ ,  $\kappa_{\text{O}_3}$ , and temperature fixed so as to consider only the consequences of the changes in  $J_{\text{O}_2}$  and  $J_{\text{O}_3}$ ).

435 The benchmark response to doubling of UV in the Chapman+2 model is shown in Fig. 9d. Doubling UV leads to minimal changes of  $\text{O}_3$  in the O-damped regime. The regime transition shifts downwards, such that peak  $\text{O}_3$  increases, and  $\text{O}_3$  also increases throughout the  $\text{O}_3$ -damped regime. The textbook paradigms and our regime transition theory make qualitatively different predictions for this response.

If the ozone layer were thought to be explained by the source/sink competition paradigm, then doubling UV would be predicted to cause no change in odd oxygen (which is dominated by  $\text{O}_3$ ) (Fig. 9a), because it rescales the photolysis of  $\text{O}_2$  and  $\text{O}_3$  by the same factor, preserving  $J_{\text{O}_2}/J_{\text{O}_3}$ .

If the ozone layer were thought to be explained by the source-controlled paradigm, then doubling UV would be predicted to increase  $\text{O}_3$  due to the increased source. The increased  $\text{O}_3$  aloft would then absorb some of the doubled UV, attenuating the UV perturbation towards the surface. This damping of the UV perturbation is a manifestation of photochemical stabilization, which is analyzed in depth in Match et al. (2024). Because of this photochemical stabilization, the steady-state  $[\text{O}_3]$  response is predicted to be top-heavy, and peak  $[\text{O}_3]$  shifts upwards and increases in magnitude.

If the ozone layer is explained by the regime transition theory, as we have argued, then doubling UV leads to a response that combines aspects of both paradigms. In the O-damped regime aloft, odd oxygen (which is dominated by ozone) does not



**Figure 9.** Response of tropical  $[O_3]$  to doubling UV at the top of the atmosphere, which perturbs photolysis rates while holding all else fixed (e.g., temperature, catalyst concentrations). (a-c) Qualitative response in each conceptual framework (described in main text) compared to a quantitative benchmark of (d) the Chapman+2 model. Only the regime transition theory correctly predicts that peak  $[O_3]$  (circles) shifts downwards in altitude and becomes larger in amplitude.

change because the the UV perturbation increases  $J_{O_2}$  and  $J_{O_3}$  by the same factor, preserving their ratio. The increase in  $J_{O_3}$  does repartition odd oxygen in favor of  $[O]$ , which strengthens the O-damping and shifts the regime transition downwards. Because the O-damped regime extends farther down, ozone has more depth over which to increase towards its peak, which correspondingly shifts downwards in altitude and increases in magnitude. Below the regime transition, in the  $O_3$ -damped regime, the enhanced UV (from the doubling) drives a top-heavy increase in ozone. Thus, the regime transition theory is the only theory to correctly explain why peak  $[O_3]$  shifts downwards and increases in magnitude.

## 455 7.2 The Chapman cycle predicts the correct altitude of peak $[O_3]$ for the wrong reason

When the scaling for ozone structure from the Chapman cycle of  $(J_{O_2}/J_{O_3})^{1/2} n_a^{3/2}$  is calculated from photolysis rates based on observed ozone, peak  $[O_3]$  is predicted to occur at 15 km (Fig. 2), which is biased far too low. However, when the Chapman cycle is calculated with photolysis rates based on its own ozone profile, it predicts an interior maximum of 24 km, quite close to the observed maximum (Fig. 3d). The success of the Chapman cycle at predicting the altitude of peak  $O_3$  underpins its reputation as the foundational model of ozone photochemistry, despite its known omission of dominant sinks. However, by comparing these two predictions, it is clear that underestimated sinks in the Chapman cycle sink cause it to overestimate ozone (by approximately a factor of two), with the consequent biases in the photolysis rates leading it to predict the correct altitude of peak  $[O_3]$  for the wrong reason.

## 8 Conclusions

465 This paper revisited the classic question of why tropical  $[O_3]$  has an interior maximum in the stratosphere around 26 km. Previous explanations in textbooks were categorized into two paradigms, each yielding a quantitative scaling for the structure of the ozone layer in terms of photolysis rates and  $[O_2]$ . Each paradigm omits the dominant sinks of ozone in their own way, and, when tested quantitatively, predict interior maxima of  $[O_3]$  that are off by 10 km in either direction (Fig. 2).

Seeking a theory of the interior maximum that is accurate to within roughly 1 km, this paper analyzed the tropical strato-  
470 spheric regime of photochemistry and transport by augmenting the Chapman cycle with linear damping of O and  $O_3$  to represent sinks from catalytic cycles and transport. From this Chapman+2 model, the textbook paradigms emerge as well-defined limits in an O-damped regime or  $O_3$ -damped regime. Yet, as before, neither regime explains the interior maximum on its own. Instead, a new theory is offered: the interior maximum of tropical stratospheric  $[O_3]$  occurs at the transition from an O-damped regime aloft to an  $O_3$ -damped regime below.

475 This regime transition is closely related to the well-known transition from a photochemically-dominated regime aloft to a transport-dominated regime below (e.g., Garcia and Solomon, 1985; Perliski et al., 1989). Here, we interpreted this regime transition in terms of competition between O-damping and  $O_3$ -damping, although it has been relatedly interpreted in terms of competition between photochemistry and transport. The regime transition emerges primarily due to increasing air density and attenuation of UV towards the surface, both of which repartition odd oxygen away from O and in favor of  $O_3$ . The fact that  
480 peak  $[O_3]$  occurs at the regime transition has not been previously noted or contextualized within a causal framework.

Under gray radiation (and other assumptions), the regime transition theory can be used to derive an analytical solution for the ozone profile (Sec. 6). Accompanying this analytical solution is an interpretive framework in which the ozone layer is conceived to begin aloft in an O-damped regime, in which ozone increases towards the surface and scales as  $(J_{O_2}/J_{O_3})n_a^3$ , before transitioning to an  $O_3$ -damped regime below in which ozone decreases towards the surface and scales as  $J_{O_2}[O_2]$ .

485 The value of a theory lies in its ability to correctly predict the response to perturbations. As shown in Section 7, the standard textbook explanations fail to even qualitatively predict the response of the ozone layer to changes in incoming UV radiation. This is because perturbations can have distinct effects in the O-damped regime, the  $O_3$ -damped regime, and on the transition between them. In the case of doubling UV, the regime transition shifts downwards, causing peak  $[O_3]$  to shift downwards and increase in magnitude. The regime transition theory could be useful for interpreting the ozone response to other perturbations.

### 490 Appendix A: Numerical details for solving the Chapman+2 model

We implement a numerical solution to the Chapman cycle by solving Eq. 7 iteratively from the top of the atmosphere downwards. At any given level, we first solve for the UV flux reaching that level, which constrains the photolysis rates  $J_{O_2}$  and  $J_{O_3}$ . These photolysis rates are then used to solve for  $O_3$  (Eq. 7), which (along with  $O_2$ ) constrains the UV flux reaching the level below. We consider the case of overhead sun. We consider damping by prescribed  $\kappa_O$  and  $\kappa_{O_3}$ , but, except as possibly  
495 accounted for by those damping rates, we do not explicitly account for advection, tropospheric chemistry, scattering, clouds, or surface reflection.

The vertical dimension is discretized into vertical levels ( $\Delta z = 100$  meters) ranging from the surface to 100 km. The idealized shortwave radiative transfer and photolysis rates are solved on a wavelength grid with 621 discretized wavelengths ranging from 180 nm to 800 nm, extending into the Chappuis bands of weak absorption. Simulated absorption in the weakly-absorbing Chappuis bands ( $\lambda > 400$  nm) is approximately  $3 \cdot 10^{-4}$  molec cm $^{-3}$  s $^{-1}$ , consistent with that reported by Nicolet (1980). Spectrally-resolved parameters are linearly interpolated to the wavelength grid. Top-of-atmosphere UV flux is calculated from the Solar Spectral Irradiance Climate Data Record (Coddington et al., 2015), based on the Naval Research Laboratory model for spectral and total irradiance and averaged over its full record from 1610-2020 (Fig. 3a). Our absorption for O $_2$  and O $_3$  are taken from Burkholder et al. (2019), where per their recommendation we use Kockarts (1976) for  $\sigma_{O_2}(\lambda < 205$  nm) (Fig. 3b). The isothermal atmosphere has a default temperature of 240 K and scale height of 7 km. Temperature-dependent parameters for reaction rates are taken from Brasseur and Solomon (2005).

## Appendix B: Simple spectral models for the Chapman cycle

The interior maximum of ozone in the Chapman cycle is of theoretical and historical significance (Chapman, 1930), yet clarity can still be gained as to how exactly this interior maximum comes about. The Chapman cycle leads to an interior maximum explained by the source/sink competition paradigm, and is in a photolytic sink regime. We clarify the role of structure in the absorption coefficients in leading to this interior maximum by using two highly-idealized simple spectral models (SSMs) (terminology after Jeevanjee and Fueglistaler, 2020), for which we replace the O $_2$  and O $_3$  absorption spectra with simple analytical functions. Once these analytical functions are embedded in the photochemical system, we elucidate how the interior maximum of the ozone layer emerges from spectral absorption features.

### B1 No interior maximum under gray radiation

The Chapman cycle can be solved explicitly in the limit of gray radiative transfer, just as in the case of the O-damped regime (Section 6.1), which also occupies a photolytic sink regime. Under gray radiation,  $J_{O_2}/J_{O_3}$  reduces to  $\sigma_{O_2}^*/\sigma_{O_3}^*$ , yielding an explicit ozone profile:

$$[O_3]_{\text{gray,Chapman}} = \left( \frac{\sigma_{O_2}^* k_2}{\sigma_{O_3}^* k_4} \right)^{1/2} C_{O_2} n_a^{3/2} \quad (\text{B1})$$

This explicit ozone profile can be integrated to yield a column ozone:

$$\chi_{O_3}(z) = \frac{2H}{3} [O_3]_{\text{gray,Chapman}} \quad (\text{B2})$$

This expression for column ozone can be substituted into explicit expressions for the photolysis rates ( $J_{O_2}$  and  $J_{O_3}$ ). The resulting gray Chapman cycle solutions are shown in Fig. A1 (top row). Because the production rate of ozone maximizes at a sweet spot in the interior of the atmosphere but  $[O_3]$  maximizes at the surface, the Chapman cycle does not generally



525 obey the source-controlled paradigm. The production rate of ozone ( $J_{O_2}[O_2]$ ) maximizes at  $\tau_{O_3} = 2/3$  even as  $O_3$  maximizes at the surface. In the source/sink competition paradigm, ozone can maximize arbitrarily far below the peak in its production rate. Lifting the ozone maximum into the interior of the atmosphere in the source/sink competition paradigm requires spectral structure.

## B2 A two-band model for peak $O_3$ in the Chapman cycle

530 Spectral structure can be incorporated with minimal complexity into our simple spectral model by adding an extra window of UV radiation, making this a two-band model. The added spectral structure is the *extension window* of ozone absorption at higher wavelengths. The extension window results because  $O_3$  can be photolyzed by photons up to 1080 nm, whereas  $O_2$  can only be photolyzed up to 240 nm. This reflects the weaker bonds of  $O_3$  compared to  $O_2$ . Thus, below 240 nm there is absorption by both  $O_2$  and  $O_3$  in an *overlap window*, whereas beyond 240 nm there is only absorption by  $O_3$  in the extension  
535 window.

We represent the extension window by extending  $O_3$  absorption to longer wavelengths where it no longer overlaps with  $O_2$  (Fig. A1d). Here, we assume that  $O_3$  has the same absorption coefficient in the overlap and extension window, and that these two windows have equal width in wavelength. This additional absorption increases the photolysis rate of  $O_3$ :

$$J_{O_2} = \mu\sigma_{O_2}^* q_{O_2}^* I_\infty \exp(-\sigma_{O_2}^* \chi_{O_2} - \sigma_{O_3}^* \chi_{O_3}) \quad (B3)$$

$$540 \quad J_{O_3} = \mu\sigma_{O_3}^* q_{O_3}^* I_\infty \exp(-\sigma_{O_2}^* \chi_{O_2} - \sigma_{O_3}^* \chi_{O_3}) + \mu\sigma_{O_3}^* I_\infty \exp(-\sigma_{O_3}^* \chi_{O_3}) \quad (B4)$$

The second term on the right-hand side of Eq. B4 is the additional photolysis in the extension window. Although  $J_{O_2}$  has the same functional form as in the gray case, note that it will not take the same values because  $\chi_{O_3}$  must refer to a self-consistent overhead column ozone profile. Plugging  $J_{O_2}$  and  $J_{O_3}$  into Eq. 11, the implicit terms again cancel leading to an explicit solution for ozone:

$$545 \quad [O_3]_{\text{Extension}}(z) = \left( \frac{\sigma_{O_2}^* k_2}{\sigma_{O_3}^* (1 + \exp(\sigma_{O_2}^* \chi_{O_2}(z))) k_4} \right)^{1/2} C_{O_2} n_a(z)^{3/2} \quad (B5)$$

This is an explicit expression for an ozone profile with an interior maximum in the Chapman cycle using the two-band SSM. The solution depends on overhead column  $O_2$  (assumed invariant). Whereas the Gray SSM had constant  $J_{O_2}/J_{O_3}$  with height, the Extension SSM has  $J_{O_2}/J_{O_3}$  decreasing towards the surface. In the limit where  $\exp(\sigma_{O_2}^* \chi_{O_2}) \gg 1$ , the maximum number density of ozone occurs at  $\tau_{O_2} = 3$ . For the parameters in Fig. A1f, this maximum occurs at 17 km. The altitude of  
550 peak  $O_3$  depends only on  $O_2$  optical depth because, with constant  $\sigma_{O_3}^*$ , absorption by  $O_2$  is what causes the photolytic source to attenuate faster than the photolytic sink.

Conceptually, in the source/sink competition paradigm, ozone maximizes in the interior of the atmosphere due to competition between the exponentially-increasing air density towards the surface and the declining ratio of the photolytic source to the photolytic sink ( $J_{O_2}/J_{O_3}$ ). The Extension SSM reveals that the photolysis rate of  $O_2$  is attenuated faster than the photolysis rate of  $O_3$  due to the joint structure of the  $O_2$  and  $O_3$  absorption coefficients, which have a region of overlapping absorption that both produces and destroys ozone and a region of extended ozone absorption that only destroys ozone. Once the overlap window saturates with  $O_2$ , its contribution to both the ozone source and sink begins to decline rapidly. Because the overlap window accounts for all of the source but only part of the sink, the sink being buttressed by contributions from the extension window, the source decreases relative to the sink.

The results from the Extension SSM suggest that the interior maximum in the photolytic sink regime is explained by the source/sink competition paradigm. These insights backstop previous explanations of ozone structure within the source/sink competition paradigm. For example, Dutsch (1968) wrote (with adapted notation), “The formation of a layer of maximum ozone content arises from the fact that below about 35 km the dissociation rate of molecular oxygen ( $J_{O_2}$ ) drops off much more rapidly than that of ozone ( $J_{O_3}$ ), mainly because of the overlap of ozone and oxygen absorption around 210 nm.” McElroy (2002) wrote that the concentration of  $O_3$  “is small at low altitudes, reflecting the *comparative absence* [emphasis added] of radiation with wavelengths sufficiently short to effect dissociation of  $O_2$ .” “Comparative” refers to the difference between the ozone production and destruction. Although these explanations are cogent examples of the source/sink competition paradigm, we reiterate that peak  $[O_3]$  is actually not explained by the Chapman cycle or its associated source/sink competition paradigm, but rather by the regime transition theory (Section 4).

## 570 Appendix C: An explicit solution to the source-controlled paradigm under gray radiation

In Section 6.3, we derived the ozone profile in the  $O_3$ -damped regime below some transition altitude  $z_t$  at which  $J_{O_2}(z_t)$  was known. Here, we derive an ozone profile for an atmosphere assumed to be everywhere in an  $O_3$ -damped regime, whose structure is therefore explained by the source-controlled paradigm. Our derivation can be generalized from that in Section 6.3 by taking  $z_t$  towards  $\infty$  and substituting  $J_{O_2}(z_t)$  as dictated by the top-of-atmosphere UV flux, i.e.,  $J_{O_2}(\infty) = \sigma_{O_2}^* \mu q_{O_2}^* I_\infty$ . This yields the following expression for ozone:

$$[O_3](z) = \frac{\sigma_{O_2}^* C_{O_2} n_{a_0} \exp(-z/H)}{\sigma_{O_3}^* ((1 + \alpha_{O_3}) \exp(\tau_{O_2}(0) \exp(-z/H)) - 1)} \quad (C1)$$

where the non-dimensional parameters  $\alpha_{O_3}$  and  $\tau_{O_2}(0)$  were defined by Eqs. 22 and 23. The values of  $\alpha_{O_3}$  must be restricted by the assumption that damping is strong enough to lead to an  $O_3$ -damped regime, which rules out values of  $\alpha_{O_3}$  below a certain threshold that can be *post hoc* verified for a given solution.

580 By differentiating Eq. C1, the ozone maximum can be found to be located at the following optical depth with respect to  $O_2$ :

$$\tau_{O_2, \max O_3} = W \left( \frac{-1}{(1 + \alpha_{O_3})e} \right) + 1 \quad (C2)$$

Eq. C2 reveals that when damping is very strong, in the limit of  $\alpha_{O_3}$  going to  $\infty$ , the interior maximum of ozone is at  $\tau_{O_2} = 1$ , i.e., at the sweet spot calculated from  $O_2$  absorption. This limit corresponds to the limit of vanishing ozone, in which  $O_2$  is the dominant absorber of UV. This recovers a textbook problem on the shape of the ozone layer from Jacob (1999) (Chapter 10),  
585 which neglects absorption by  $O_3$ . However, as damping weakens to the point that  $O_3$  increases enough to become the dominant absorber, while still ensuring that the damping is strong enough to be in the  $O_3$ -damped regime, absorption by ozone suppresses the production rate at lower altitudes and shifts the interior maximum in ozone production (and ozone itself) towards higher altitudes.


Fig. A2 shows how the theoretical scaling compares with numerical solutions to the Chapman cycle under gray radiation  
590 with  $O_3$  damping. The theoretical scaling correctly captures that, for strong damping, the ozone maximum approaches  $\tau_{O_2} = 1$ , which is the gray  $O_2$ -only limit. As damping is reduced, the theoretical scaling correctly captures that the interior maximum shifts upwards as absorption by ozone aloft reduces the ozone production rate at lower altitudes. However, further reductions in damping lead to the violation of the underlying assumptions of the theoretical scaling, namely that ozone is everywhere in the  $O_3$ -damped regime. Instead, the Chapman cycle sink of ozone can dominate in the upper atmosphere, leading to a photolytic  
595 sink regime aloft unaccounted-for by this theory. Thus, the theory performs well in its range of applicability, but does not explain the observed ozone maximum on Earth.

*Code and data availability.* Chapman+2 Photochemical-Transport Model described in Section 2 is published on Zenodo at <https://doi.org/10.5281/zenodo.13412268>.

*Author contributions.* Authors' contributions: AM and EPG acquired funding; AM, EPG, and SF conceptualized research; AM performed  
600 formal analysis; AM wrote original draft; EPG and SF reviewed and edited paper.

*Competing interests.* The authors declare that they have no conflict of interest.


*Acknowledgements.* The authors thank editor Jens-Uwe Grooß and two anonymous reviewers for their valuable feedback. We also thank Daniel Cariolle for sharing the latest version (v2.9) of his linear ozone model. A.M. acknowledges constructive discussions with Benjamin Schaffer, Nadir Jeevanjee, Nathaniel Tarshish, and at the Princeton Center for Theoretical Science workshop *From Spectroscopy to Climate*.  
605 This work was supported by the National Science Foundation under Award No. 2120717 and OAC-2004572, and by Schmidt Sciences, a philanthropic initiative founded by Eric and Wendy Schmidt, as part of the Virtual Earth System Research Institute (VESRI).



**Figure A1.** Chapman cycle photochemical equilibrium in simple spectral models of the tropical ozone layer (first two rows) and the full spectral model (bottom row). (Left column) Absorption coefficients in each model (solid) compared to in reality (transparent). (Middle column)  $\text{O}_2$  and  $\text{O}_3$  photolysis rates. (Right column) Ozone number density.

## References

- Bates, D. R. and Nicolet, M.: The Photochemistry of Atmospheric Water Vapor, *Journal of Geophysical Research* (1896-1977), 55, 301–327, <https://doi.org/10.1029/JZ055i003p00301>, 1950.
- 610 Brasseur, G., Hitchman, M. H., Walters, S., Dymek, M., Falise, E., and Pirre, M.: An Interactive Chemical Dynamical Radiative Two-Dimensional Model of the Middle Atmosphere, *Journal of Geophysical Research: Atmospheres*, 95, 5639–5655, <https://doi.org/10.1029/JD095iD05p05639>, 1990.
- Brasseur, G. P. and Jacob, D. J.: Modeling of Atmospheric Chemistry, *Modeling of Atmospheric Chemistry*, pp. 1–606, <https://doi.org/10.1017/9781316544754>, 2017.



**Figure A2.** Altitude of peak tropical  $[O_3]$  under gray radiation with absorption by  $O_2$  and  $O_3$  and varying damping  $\kappa_{O_3}^*$  that modulates the non-dimensional  $O_3$ -damping parameter  $\alpha_{O_3}$ . Comparison of numerical simulations (black) with analytical theory (red, Eq. C2). The theory reproduces the strongly damped limit where peak  $[O_3]$  occurs at  $\tau_{O_2} = 1$ . As damping is weakened, absorption by ozone aloft weakens the ozone source below and shifts peak  $[O_3]$  upwards. As damping is further weakened, the theoretical assumption of a non-photolytic sink regime breaks down as the Chapman cycle sink becomes important aloft, degrading the applicability of the strongly-damped theory.

- 615 Brasseur, G. P. and Solomon, S.: *Aeronomy of the Middle Atmosphere: Chemistry and Physics of the Stratosphere and Mesosphere*, Springer, Dordrecht, Netherlands, 2005.
- Burkholder, J. B., Sander, S. P., Abbatt, J., Barker, J. R., Cappa, C., Crouse, J. D., Dibble, T. S., Huie, R. E., Kolb, C. E., Kurylo, M. J., Orkin, V. L., Percival, C. J., Wilmouth, D. M., and Wine, P. H.: *Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies*, Tech. rep., Jet Propulsion Laboratory, Pasadena, CA, 2019.
- 620 Chapman, S.: A Theory of Upper Atmospheric Ozone, *Memoirs of the Royal Meteorological Society*, III, 103–125, 1930.
- Coddington, O., Lean, J., Lindholm, D., Pilewskie, P., and Snow, M.: *NOAA Climate Data Record (CDR) of Solar Spectral Irradiance (SSI)*, Version 2.1, <https://doi.org/10.7289/V53776SW>, 2015.
- Craig, R. A.: *The Upper Atmosphere: Meteorology and Physics*, Academic Press, ISBN 978-0-12-194850-4, 1965.
- Crutzen, P. J.: The Influence of Nitrogen Oxides on the Atmospheric Ozone Content, *Quarterly Journal of the Royal Meteorological Society*, 625 96, 320–325, <https://doi.org/10.1002/qj.49709640815>, 1970.
- Davis, S. M., Rosenlof, K. H., Hassler, B., Hurst, D. F., Read, W. G., Vömel, H., Selkirk, H., Fujiwara, M., and Damadeo, R.: The Stratospheric Water and Ozone Satellite Homogenized (SWOOSH) Database: A Long-Term Database for Climate Studies, *Earth System Science Data*, 8, 461–490, <https://doi.org/10.5194/essd-8-461-2016>, 2016.
- Dobson, G. M. B.: Origin and Distribution of the Polyatomic Molecules in the Atmosphere, *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 236, 187–193, <https://doi.org/10.1098/rspa.1956.0127>, 1956.
- 630 Dobson, G. M. B.: *Exploring the Atmosphere*, by G.M.B. Dobson., Clarendon Press, Oxford, 1963.

- Dutsch, H. U.: *The Photochemistry of Stratospheric Ozone*, Royal Meteorological Society, 94, 483–497, 1968.
- Garcia, R. R. and Solomon, S.: The Effect of Breaking Gravity Waves on the Dynamics and Chemical Composition of the Mesosphere and Lower Thermosphere, *Journal of Geophysical Research*, 90, 3850, <https://doi.org/10.1029/JD090iD02p03850>, 1985.
- 635 Hartley, W. N.: XXI.—On the Absorption of Solar Rays by Atmospheric Ozone, *Journal of the Chemical Society, Transactions*, 39, 111–128, <https://doi.org/10.1039/CT8813900111>, 1881.
- Jacob, D.: *Introduction to Atmospheric Chemistry*, Princeton University Press, 1999.
- Jeevanjee, N. and Fueglistaler, S.: Simple Spectral Models for Atmospheric Radiative Cooling, *Journal of the Atmospheric Sciences*, 77, 479–497, <https://doi.org/10.1175/JAS-D-18-0347.1>, 2020.
- 640 Jeevanjee, N., Seeley, J. T., Paynter, D., and Fueglistaler, S.: An Analytical Model for Spatially Varying Clear-Sky CO<sub>2</sub> Forcing, *Journal of Climate*, 34, 9463–9480, <https://doi.org/10.1175/JCLI-D-19-0756.1>, 2021.
- Keeble, J., Hassler, B., Banerjee, A., Checa-Garcia, R., Chiodo, G., Davis, S., Eyring, V., Griffiths, P. T., Morgenstern, O., Nowack, P., Zeng, G., Zhang, J., Bodeker, G., Burrows, S., Cameron-Smith, P., Cugnet, D., Danek, C., Deushi, M., Horowitz, L. W., Kubin, A., Li, L., Lohmann, G., Michou, M., Mills, M. J., Nabat, P., Olivie, D., Park, S., Seland, Ø., Stoll, J., Wieners, K. H., and Wu, T.: Evaluating Stratospheric Ozone and Water Vapour Changes in CMIP6 Models from 1850 to 2100, *Atmospheric Chemistry and Physics*, 21, 5015–5061, <https://doi.org/10.5194/ACP-21-5015-2021>, 2021.
- 645 Kockarts, G.: Absorption and Photodissociation in the Schumann-Runge Bands of Molecular Oxygen in the Terrestrial Atmosphere, *Planetary and Space Science*, 24, 589–604, [https://doi.org/10.1016/0032-0633\(76\)90137-9](https://doi.org/10.1016/0032-0633(76)90137-9), 1976.
- Kump, L. R., Kasting, J. F., and Crane, R. G.: *The Earth System*, Pearson Education, ISBN 978-0-321-73328-3, 2011.
- 650 Liou, K.: *An Introduction to Atmospheric Radiation*, Academic Press, ISBN 978-0-12-451451-5, 2002.
- Match, A. and Gerber, E. P.: Tropospheric Expansion Under Global Warming Reduces Tropical Lower Stratospheric Ozone, *Geophysical Research Letters*, 49, e2022GL099463, <https://doi.org/10.1029/2022GL099463>, 2022.
- Match, A., Gerber, E. P., and Fueglistaler, S.: Beyond Self-Healing: Stabilizing and Destabilizing Photochemical Adjustment of the Ozone Layer, *Atmospheric Chemistry and Physics*, 24, 10305–10322, <https://doi.org/10.5194/acp-24-10305-2024>, 2024.
- 655 McElroy, M. B.: *The Atmospheric Environment*, <https://doi.org/10.1515/9780691234663/HTML>, 2002.
- Neu, J. L. and Plumb, R. A.: Age of Air in a “Leaky Pipe” Model of Stratospheric Transport, *Journal of Geophysical Research*, 104, 19243, <https://doi.org/10.1029/1999JD900251>, 1999.
- Nicolet, M.: Solar UV Radiation and Its Absorption in the Mesosphere and Stratosphere, *pure and applied geophysics*, 118, 3–19, <https://doi.org/10.1007/BF01586443>, 1980.
- 660 Perliski, L. M., Solomon, S., and London, J.: On the Interpretation of Seasonal Variations of Stratospheric Ozone, *Planetary and Space Science*, 37, 1527–1538, [https://doi.org/10.1016/0032-0633\(89\)90143-8](https://doi.org/10.1016/0032-0633(89)90143-8), 1989.
- Pierrehumbert, R. T.: Infrared Radiation and Planetary Temperature, *Physics Today*, 64, 33–38, <https://doi.org/10.1063/1.3541943>, 2011.
- Romps, D. M., Seeley, J. T., and Edman, J. P.: Why the Forcing from Carbon Dioxide Scales as the Logarithm of Its Concentration, *Journal of Climate*, 35, 4027–4047, <https://doi.org/10.1175/JCLI-D-21-0275.1>, 2022.
- 665 Seinfeld, J. H. and Pandis, S. N.: *Atmospheric Chemistry and Physics: From Air Pollution to Climate Change*, 3rd Edition | Wiley, Wiley, 2016.
- Visconti, G.: *Fundamentals of Physics and Chemistry of the Atmosphere*, Springer International Publishing, second edn., 2016.